

UG-FT-214/07
 CAFPE-84/07
 FFUOV-07/02
 hep-th/0701151
 January 2007

Adding magnetic flux to the baryon vertex

Bert Janssen^{a,1}, Yolanda Lozano^{b,2} and Diego Rodríguez-Gómez^{c,3}

*^aDepartamento de Física Teórica y del Cosmos and
 Centro Andaluz de Física de Partículas Elementales
 Universidad de Granada, 18071 Granada, Spain*

*^bDepartamento de Física, Universidad de Oviedo,
 Avda. Calvo Sotelo 18, 33007 Oviedo, Spain*

*^c Department of Physics, Princeton University,
 Princeton, NJ 08540, U.S.A.*

ABSTRACT

We generalise the baryon vertex configuration of AdS/CFT by adding magnetic field on its world-volume, dissolving D-string charge. A careful analysis of the configuration shows that there is an upper bound on the number of dissolved strings. We provide a microscopical description of this configuration in terms of a dielectric effect for the dissolved strings.⁴

¹E-mail address: bjanssen@ugr.es

²E-mail address: yolanda@string1.ciencias.uniovi.es

³E-mail address: drodrigu@princeton.edu

⁴Talk given by B.J. at the RTN Network Meeting in Napoli, in Octobre 2006.

1 Introduction

The AdS/CFT correspondence conjectures a remarkable duality between strings moving in an $AdS_5 \times S^5$ background with N units of 4-form R-R flux and $D = 4$, $\mathcal{N} = 4$ SYM in flat space with gauge group $SU(N)$ [1]. In particular, the conjecture states that quark and anti-quarks in the $SU(N)$ gauge theory manifest themselves in $AdS_5 \times S^5$ as the endpoints of open strings ending on the boundary of the AdS_5 .

It is clear then that one can form a $q\bar{q}$ state with a single U-shaped string coming from the boundary, entering into the bulk, with the apex at a distance u_0 from the centre of the AdS_5 and then going back to the boundary. This configuration is seen on the SYM side as a $q\bar{q}$ pair (a meson), whose energy is computed by means of a Wilson loop, while on the gravity side the energy is computed by minimising the worldsheet area of the string ending on the loop [2, 3].

One would expect that, besides the meson, there should be a mechanism to form a baryon configuration, *i.e.* an anti-symmetric, colourless, finite-energy bound state of N non-dynamical quarks. In [4] the gravitational dual of such bound state of quarks was found in terms of a D5-brane wrapping the S^5 part of the $AdS_5 \times S^5$ background. On this D5-brane there are N F-strings attached, stretching from the D5-brane to the boundary of AdS_5 , whose endpoints are regarded on the SYM side as a bound state of N quarks. It can be shown [4] that the associated wave function satisfies the required symmetry properties.

The best way to see that the above configuration is indeed the baryon vertex is by looking at the Chern-Simons term of the D5-brane, wrapped around the S^5 . Since in $AdS_5 \times S^5$ there is no 6-form R-R potential to which the probe brane can couple, the only non-zero term in the Chern-Simons action is the 4-form R-R field $C^{(4)}$ coupled to the BI field strength $F = dA$. In our specific setting, the only non-zero contribution is that of the coupling of the magnetic part of $C^{(4)}$ to the electric component of F . Integrating by parts, this term can then be rewritten as

$$S_{CS} = -T_5 \int_{\mathbb{R} \times S^5} P[C^{(4)}] \wedge F = T_5 \int_{\mathbb{R} \times S^5} P[G^{(5)}] \wedge A, \quad (1.1)$$

where $G^{(5)} = dC^{(4)}$ is the R-R 5-form field strength. In $AdS_5 \times S^5$ we have that $\int_{S^5} G^{(5)} = 4\pi^2 N$ (in units where $2\pi l_s^2 = 1$), such that with the Ansatz $A = A_t(t)dt$, it is clear that the coupling (1.1) factorises as

$$S_{CS} = T_5 \int_{S^5} G^{(5)} \int dt A_t = NT_1 \int dt A_t, \quad (1.2)$$

where we have taken into account that the tension of the D5-brane and the tension of a string are related by $4\pi^2 T_5 = T_1$. Therefore, the coupling (1.1) is inducing N units of electric BI charge on the D5-brane. However we have to check whether the Ansatz for A is consistent with the equations of motion of the total D5-brane system. The equation of motion of A is given by $NT_1 = 0$, such that the above construction is only consistent if the total electric BI charge on the D5-brane is zero. Yet, there is a way of inducing a non-zero BI electric charge in the worldvolume of the D5-brane, by cancelling this charge with the charge induced by the endpoints of N open fundamental strings (with appropriate orientation) stretching between the D5-brane and the boundary of the AdS space. The full dynamics of the configuration is therefore not only described by the D5-brane action, but also by the action for the open strings, consisting of N copies of the Nambu-Goto action S_{F1} and a boundary term contribution $T_1 \int A_t dt$ from the endpoints. Note that the contribution from the open string endpoints cancels exactly the Chern-Simons term in the D5-brane action, such that the total system is described by [5]

$$S_{\text{total}} = S_{DBI} + NT_1 \int dt A_t + NS_{F1} - NT_1 \int dt A_t = S_{DBI} + NS_{F1}. \quad (1.3)$$

The configuration that we have just described is the so-called baryon vertex. Since the N F-strings, stretching from the D5-brane all the way to the AdS boundary, have the same orientation, the dual configuration on the CFT side corresponds to the bound state of N (anti)quarks, which is gauge invariant and antisymmetric under the interchange of any two quarks [4].

We will show that the baryon vertex can be generalised by adding a new quantum number, representing magnetic flux. The key point is to realise that S^5 can be seen as an S^1 bundle over CP^2 . The S^1 fibre is a non-trivial $U(1)$ gauge bundle on the CP^2 base, and this allows to switch on a magnetic BI field on the worldvolume of the D5-brane, proportional to the curvature tensor of the fibre connection. We will show that the magnetic flux leads to surprising dynamics and a complementary description of the baryon vertex as a dielectric effect. The results mentioned in this letter are discussed in detail in [6].

2 A generalisation of the baryon vertex

It is well known that S^5 can be regarded as a $U(1)$ fibre over CP^2 with a non-trivial fibre connection B . In these S^5 fibre coordinates the $AdS_5 \times S^5$ background then reads

$$\begin{aligned} ds^2 &= \frac{u^2}{L^2} \eta_{ab} dx^a dx^b + \frac{L^2}{u^2} du^2 + L^2 \left((d\chi - B)^2 + ds_{CP^2}^2 \right), \\ C_{abcd} &= L^{-4} u^4 \epsilon_{abcd}, \quad C_{\varphi_2 \varphi_3 \varphi_4 \chi} = \frac{1}{8} L^4 \sin^4 \varphi_1 \sin \varphi_2, \end{aligned} \quad (2.4)$$

where $ds_{CP^2}^2$ stands for the Fubini-Study metric on CP^2 (with coordinates φ_i) and χ is taken along the $U(1)$ fibre. The fibre connection B satisfies the following properties [7]

$$dB = \star(dB), \quad \int_{CP^2} dB \wedge dB = 4\pi^2, \quad (2.5)$$

where the Hodge star is taken with respect to the Fubini-Study metric on CP^2 . In particular, one sees that B induces a non-trivial instanton number in CP^2 [7, 8]. In these coordinates the baryon vertex consists of the D5-brane wrapped around the S^5 and the fundamental strings laying in the u -direction of AdS_5 [3]. Besides the electric components of the BI field strength of the previous section, one could think of turning on also magnetic components. Due to (2.5), it is natural to take the magnetic components living in the CP^2 and proportional to the curvature tensor of the $U(1)$ fibre connection, $F = 2n dB$. With this Ansatz F satisfies the same properties (2.5) as the fibre connection, namely it is selfdual and

$$\int_{CP^2} F \wedge F = 8\pi^2 n^2. \quad (2.6)$$

With this choice for the BI field strength it is clear that there are no other couplings in the Chern-Simons action besides the ones we already considered in (1.1). The Born-Infeld action however is given by

$$S_{DBI} = -T_5 \int d^6 \xi \frac{u}{L} \sqrt{\det(g_{\alpha\beta} + F_{\alpha\beta})} = -T_5 \int d^6 \xi u \sqrt{g_{S^5}} \left(L^4 + 2F_{\alpha\beta} F^{\alpha\beta} \right), \quad (2.7)$$

where we have used that F is selfdual, such that the determinant under the square root is a perfect square. The Ansatz for F is consistent with the action (2.7), as is reflected in the fact that the equations of motion for the magnetic components of F are given by $dF = 0$. Finally, substituting the expression for F in the action and integrating over the S^5 directions we obtain the following expression for the energy of the spherical D5-brane:

$$E_{D5} = 8\pi^3 T_5 u \left(n^2 + \frac{L^4}{8} \right). \quad (2.8)$$

Note that this energy consists of two parts: one contribution from the tension of the 5-brane wrapped around the S^5 , proportional to L^4 and one from the magnetic flux of the BI vector, proportional to n^2 .

The magnetic components of F induce a non-zero instanton number n^2 on the worldvolume of the D5-brane, since integrating the Chern-Simons coupling over the CP^2 directions, one obtains

$$S_{CS} = \frac{1}{2}T_5 \int_{\mathbb{R} \times S^5} P[C^{(2)}] \wedge F \wedge F = n^2 T_1 \int_{\mathbb{R} \times S^1} P[C^{(2)}], \quad (2.9)$$

where we have used again that $T_1 = 4\pi^2 T_5$. Even though in $AdS_5 \times S^5$ $C^{(2)}$ is zero, this coupling indicates that the magnetic flux is inducing n^2 D-string charge in the configuration. These strings are wound around the fibre direction χ and dissolved in the CP^2 . Note that the total energy of the configuration (2.8) is the sum of the energy of the D5 and the D1-branes, which hints at that we are dealing with a threshold BPS bound state.

3 A bound on the magnetic flux

It was argued in [5] that in order to analyse the stability of the baryon vertex in the u -direction, one has to consider the influence of the external F-strings. The energy E of the baryon vertex is inversely proportional to the distance ℓ between the quarks, with a negative proportionality constant, such that the baryon vertex is indeed stable under perturbations in u . In this subsection we will perform the same calculation in [5], but taking into account the effect of the non-zero magnetic flux on the D5-brane.

The equations of motion of F-strings in the baryon vertex are twofold: the bulk equation of motion for the strings and the boundary equation of motion, which contains a term coming from the D5-brane. Parametrising the worldsheet of the F-strings by $\{t, x\}$ and the position in AdS by $u = u(x)$, we obtain

$$\frac{u^4}{\sqrt{(u')^2 + \frac{u^4}{L^4}}} = \text{const}, \quad \frac{u'_0}{\sqrt{(u'_0)^2 + \frac{u_0^4}{L^4}}} = \frac{\pi L^4}{4N} \left(1 + \frac{8n^2}{L^4}\right), \quad (3.10)$$

for the bulk and the boundary respectively, with u_0 the position of the baryon vertex in the holographic direction and $u'_0 = u'(u_0)$. The equations (3.10) can be combined into a single one,

$$\frac{u^4}{\sqrt{(u')^2 + \frac{u^4}{L^4}}} = \beta u_0^2 L^2, \quad \text{with } \beta^2 = 1 - \frac{1}{16} \left(1 + \frac{8\pi n^2}{N}\right)^2. \quad (3.11)$$

In the absence of magnetic BI flux on the worldvolume, $\beta = \sqrt{15/16}$, as in [5]. However, in general for non-zero n^2 , we have to make sure that β is real (as u is real), which implies that $n^2 \leq 3N/8\pi$. Surprisingly, we find that there is a bound on the number of D-strings that can be dissolved in the configuration, which depends on the number of D3-branes that source the background.

Integrating the equation of motion, we find that the size ℓ and the energy E of the baryon are given by

$$\ell = \frac{L^2}{u_0} \int_1^\infty dy \frac{\beta}{y^2 \sqrt{y^4 - \beta^2}}, \quad E = T_1 u_0 \left\{ \int_1^\infty dy \left[\frac{y^2}{\sqrt{y^4 - \beta^2}} - 1 \right] - 1 \right\}, \quad (3.12)$$

with $y = u/u_0$. These integrals can be solved in terms of hypergeometric functions [5]. In Figure 1 we have plotted the radius and the energy of the baryon as a function of n^2/N . The plots reveal that the size of the baryon vertex goes to zero as we approach the bound on n^2 , making it impossible to continue beyond the bound. The expression for the energy has the same form as the expression in [5] and indeed takes the same value for $n = 0$. In particular, the dependence on $\sqrt{g^2 N}$ and on u_0 is unaltered, as expected by conformal invariance. Notice that also here the energy of the configuration is only well defined for n^2 inside the allowed interval.

The fact that we find a bound on the number of dissolved D1-branes due to the dynamics of the F-strings, is quite surprising. The bound is probably related to the stringy exclusion principle




Figure 1: Radius ℓ (in units of L^2/u_0) and the energy E (in units of u_0) of the baryon vertex as a function of n^2/N .

of [9]. Our configuration carries a non-zero winding number in the fibre direction of the S^5 , which in terms of the dual field theory will manifest itself as a specific charge of the $SU(3)$ R-symmetry group. As these charges are bounded due to conformal invariance, one expects to find a bound on the magnetic flux.

4 A microscopical description in terms of D1-branes

The fact that the magnetic flux induces D1-brane charge through the coupling (2.9), suggests a close analogy with the dielectric effect described in [10, 11]. In this section we will show that it is indeed possible to give an alternative, microscopic description of the baryon vertex, in terms of a fuzzy spherical D5-brane built up out of dielectrically expanded D1-branes.

The action describing the dynamics of n^2 coinciding D1-branes is the non-Abelian action given in [11], which for the $AdS_5 \times S^5$ background reduces to the form

$$S_{n^2 D1} = -T_1 \int d^2 \xi \text{STr} \left\{ \sqrt{\det \left(P[g_{\mu\nu} + g_{\mu i}(Q^{-1} - \delta)^i_j g^{jk} g_{k\nu}] \det Q \right)} \right\} \quad (4.13)$$

where $g_{\mu\nu}$ is the metric in $AdS_5 \times S^5$ and $Q^i_j = \delta^i_j + i[X^i, X^k]g_{kj}$. Inspired by the coupling (2.9) in the D5-brane calculation we wind the D-strings around the $U(1)$ fibre direction χ and let them expand into the CP^2 . In this way we obtain a fuzzy version of the S^5 as an Abelian $U(1)$ fibre over a fuzzy CP^2 [12].

A fuzzy version of CP^2 (see for example [13]) can be obtained by the set of matrix coordinates

$$X^i = \frac{T^i}{\sqrt{(2n^2 - 2)/3}}, \quad [X^i, X^j] = \frac{if^{ijk}}{\sqrt{(2n^2 - 2)/3}} X^k, \quad (4.14)$$

with T^i the generators of $SU(3)$ in the n^2 -dimensional irreducible representations $(k, 0)$ or $(0, k)$, with $n^2 = (k+1)(k+2)/2$ (see [12, 13] for more details) and with f^{ijk} the structure constant of $SU(3)$ in the algebra of the Gell-Mann matrices $[\lambda^i, \lambda^j] = 2if^{ijk}\lambda^k$.

Substituting the non-commutative Ansatz above in the action (4.13) we find

$$S_{n^2 D1} = -T_1 \int dt d\chi u \text{STr} \left\{ \mathbb{1} + \frac{L^4}{4(2n^2 - 2)} \mathbb{1} \right\} = -2\pi n^2 T_1 \int dt u \left(1 + \frac{L^4}{8(n^2 - 1)} \right), \quad (4.15)$$

since remarkably $\det Q = (\mathbb{1} + \frac{L^4}{4(2n^2 - 2)} \mathbb{1})^2$ in the large n^2 limit. It is to be emphasised that the fact that the $\det Q$ is a perfect square is the microscopical analogous of the perfect square that we obtained for the DBI action in the macroscopic case. The energy of the n^2 expanded D1-branes is then given by

$$E_{n^2 D1} = 2\pi u T_1 \left(n^2 + \frac{n^2 L^4}{8(n^2 - 1)} \right). \quad (4.16)$$

Taking into account that the tensions of the D1- and the D5-brane are related by $T_1 = 4\pi^2 T_5$, it is easy to see that in the limit where the number of D1-branes $n^2 \rightarrow \infty$, the above expression reduces to the energy of the macroscopic D5-brane, given by (2.8).

It is also possible to see the external F-strings in the microscopical description, by looking at the CS action for coincident D-strings. The coupling with the R-R 4-form and the $U(n^2)$ BI field strength $\mathcal{F} = d\mathcal{A} + [\mathcal{A}, \mathcal{A}]$ is, after partial integration and in the gauge $\mathcal{A}_\chi = 0$, given by

$$S_{CS} = -\frac{T_1}{4} \int dt d\chi \text{STr} \left\{ [X^i, X^j] [X^k, X^l] G_{\chi^{ijkl}}^{(5)} \mathcal{A}_t \right\} = \frac{L^4 T_1}{2(n^2 - 1)} \int dt d\chi \text{STr} \left\{ \mathcal{A}_t \right\}, \quad (4.17)$$

where we have used that in the non-commutative coordinates introduced in (4.14), $G^{(5)}$ is given by [12] $G_{\chi^{ijkl}}^{(5)} = L^4 f_{[ij}^m f_{kl]}^n X^m X^n$. In analogy with the Abelian case, we can take as an Ansatz for the BI vector, $\mathcal{A} = A_t(t) \mathbb{1} dt$, such that after integrating over χ , we find finally that

$$S_{CS} = \frac{n^2}{n^2 - 1} N T_1 \int dt A_t. \quad (4.18)$$

The coupling (4.17) is therefore inducing, in the large n^2 limit, N BI charges in the configuration. These charges have to be cancelled by N fundamental strings ending on the D1-brane system. The dielectric coupling to $C^{(4)}$ in (4.17) will then take care that these strings are expanded over the full S^5 .

References

- [1] J. Maldacena, Adv. Theor. Math. Phys. 2 (1998) 231, hep-th/9711200.
- [2] J. Maldacena, Phys. Rev. Lett. 80 (1998) 4859, hep-th/9803002.
- [3] S. Rey, J. Yee, Eur. Phys. J. C22 (2001) 379, hep-th/9803001.
- [4] E. Witten, JHEP 9807 (1998) 006, hep-th/9805112.
- [5] A. Brandhuber, N. Itzhaki, J. Sonnenschein, S. Yankielowicz, JHEP 9807 (1998) 020, hep-th/9806158.
- [6] B. Janssen, Y. Lozano, D. Rodríguez-Gómez, JHEP 0611 (2006) 082, hep-th/0606264.
- [7] A. Trautman, Int. J. Theor. Phys. 16 (1977) 561.
- [8] S.W. Hawking, C.N. Pope, Phys. Lett. B73 (1978) 42.
- [9] J. Maldacena, A. Strominger, JHEP 9812 (1998) 005, hep-th/9804085.
- [10] R. Emparan, Phys. Lett. B423 (1998) 71, hep-th/9711106.
- [11] R. Myers, JHEP 9912 (1999) 022, hep-th/9910053.
- [12] B. Janssen, Y. Lozano, D. Rodríguez-Gómez, Nucl. Phys. B712 (2005) 371, hep-th/0411181.
- [13] G. Alexanian, A.P. Balachandran, G. Immirzi, B. Ydri, J. Geom. Phys. 42 (2002) 28, hep-th/0103023.